

Higher Loop Effects on Unification via Kaluza-Klein ThresholdsZurab Kakushadze^{1,2,3*} and Tomasz R. Taylor^{2†}¹*Jefferson Laboratory of Physics, Harvard University, Cambridge, MA 02138*²*Department of Physics, Northeastern University, Boston, MA 02115*³*Enrico Fermi Institute, University of Chicago, Chicago, IL 60637*

(May 18, 1999)

Abstract

We discuss higher loop corrections to gauge coupling renormalization in the context of gauge coupling unification via Kaluza-Klein thresholds. We show that in the case of $\mathcal{N} = 1$ supersymmetric compactifications the one-loop threshold contributions are dominant, while the higher loop correction are subleading. This is due to the fact that at heavy Kaluza-Klein levels the spectrum as well as the interactions are $\mathcal{N} = 2$ supersymmetric. In particular, we give two different arguments leading to this result - one is field theoretic, while the second one utilizes the power of string perturbation techniques. To illustrate our discussions we perform explicit two-loop computations of various corrections to gauge couplings within this framework. We also remark on phenomenological applications of our discussions in the context of TeV-scale brane world.

Typeset using REVTeX

*E-mail: zurab@string.harvard.edu

†E-mail: taylor@neu.edu

I. INTRODUCTION

D-branes [1] are likely to play an important role in describing nature. In particular, the Standard Model gauge and matter fields may live inside of $p \leq 9$ spatial dimensional p -branes, while gravity lives in a larger (10 or 11) dimensional bulk of the space-time. This “Brane World” picture¹ *a priori* appears to be a viable scenario, and, based on considerations of gauge and gravitational coupling unification, dilaton stabilization and weakness of the Standard Model gauge couplings, in [11] it was actually argued to be a likely description of nature. In particular, these phenomenological constraints seem to be embeddable in the brane world scenario (with the Standard Model fields living on branes with $3 < p < 9$), which therefore might provide a coherent picture for describing our universe [11]². This is largely due to a much higher degree of flexibility of the brane world scenario compared with, say, the old perturbative heterotic framework.

As an example consider the gauge and gravitational couplings which in string theory are expected to unify (up to an order one factor due to various thresholds [16,17]) at the string scale $M_s = 1/\sqrt{\alpha'}$. In the brane world scenario *a priori* the string scale can be anywhere between the electroweak scale M_{ew} and the Planck scale $M_P = 1/\sqrt{G_N}$ (where G_N is the Newton’s constant). If we assume that the bulk is ten dimensional, then the four dimensional gauge and gravitational couplings scale as³ $\alpha \sim g_s/V_{p-3}M_s^{p-3}$ respectively $G_N \sim g_s^2/V_{p-3}V_{9-p}M_s^8$, where g_s is the string coupling, and V_{p-3} and V_{9-p} are the compactification volumes inside and transverse to the p -branes, respectively. For $3 < p < 9$ there are two *a priori* independent volume factors, and, for the fixed gauge coupling α (at the unification, that is, string scale) and four dimensional Planck scale M_P , the string scale is not determined. This observation was used in [2] to argue that the gauge and gravitational coupling unification problem⁴ can be ameliorated in this context by lowering the string scale M_s down to the GUT scale $M_{GUT} \approx 2 \times 10^{16}$ GeV [20]⁵. In [3] it was noticed that M_s can be further lowered all the way down to TeV.

In fact, in the brane world picture *a priori* the string scale can be as low as desired as long as it does not directly contradict current experimental data. In [4] it was proposed

¹For recent developments, see, *e.g.*, [2–12]. Large radius compactifications were originally discussed in [13].

²The brane world picture in the effective field theory context was discussed in [14,15].

³For illustrative purposes here we are using the corresponding tree-level relations in Type I (or Type I’) theory.

⁴For a review of the gauge and gravitational coupling unification problem in the perturbative heterotic string context, see, *e.g.*, [18], and references therein. In the Type I context the discussions on this issue can be found in [19,11].

⁵By the GUT scale here we mean the usual scale of gauge coupling unification in the MSSM obtained by extrapolating the LEP data in the assumption of the standard “desert” scenario.

that M_s as well as⁶ the fundamental (10 or 11 dimensional) Planck scale can be around TeV. The observed weakness of the four dimensional gravitational coupling then requires the presence of at least two large ($\gg 1/M_s$) compact directions (which are transverse to the branes on which the Standard Model fields are localized). A general discussion of possible brane world embeddings of such a scenario was given in [7,8,11]. In [10] various non-trivial phenomenological issues were discussed in the context of the TeV-scale brane world scenario, and it was argued that this possibility does not appear to be automatically ruled out⁷.

However, in such a scenario, as well as in any scenario where $M_s \ll M_{GUT}$, the gauge coupling unification at M_s would have to arise in a way drastically different from the usual MSSM unification which occurs with a remarkable precision [20]. It is then desirable to have a mechanism in the TeV-scale brane world scenario for lowering the unification scale. Moreover, it would also be necessary to find a concrete extension of the MSSM (where this new mechanism is realized) such that the unification prediction is just as precise as in the MSSM (at least at one loop). In fact, one could also require that such an extension *explain* why couplings unify in the MSSM at all, that is, why the unification in the MSSM is *not* just an “accident” assuming that the TeV-scale brane world scenario has the pretense of replacing the old framework.

In the brane world picture there appears to exist a mechanism [5] for lowering the unification scale. Thus, let the “size” R of the compact dimensions inside of the p -brane (where $p > 3$) be somewhat large compared with $1/M_s$. Then the evolution of the gauge couplings above the Kaluza-Klein (KK) threshold $1/R$ is no longer logarithmic but power-like [22]. This observation was used in [5] to argue that the gauge coupling unification might occur at a scale (which in the brane world context would be identified with the string scale) much lower than M_{GUT} .

In [12] a TeV-scale Supersymmetric Standard Model (TSSM) was proposed. The gauge coupling unification in the TSSM indeed occurs via such a higher dimensional mechanism. Moreover, the unification in the TSSM is as precise (at one loop) as in the MSSM, and occurs in the TeV range⁸. The TSSM also explains why the unification in the MSSM is not an accident - if the TSSM is indeed (a part of) the correct description of nature above the electroweak scale, then the gauge coupling unification in the MSSM is explained by the current lack of data which leads to the standard “desert” assumption. Moreover, as was pointed out in [12], after a rather systematic search the TSSM is the only (simple) solution for the constraints guaranteeing that the gauge couplings unify as precisely (at one loop) as in the MSSM.

⁶Note that the string scale M_s cannot be too much lower than the fundamental Planck scale or else the string coupling g_s as well as all the gauge couplings would come out too small contradicting the experimental data.

⁷TeV-scale compactifications were studied in [32] in the context of supersymmetry breaking.

⁸By the TeV range we do *not* necessarily mean that $M_s \sim 1$ TeV. In fact, as was argued in [12], the gauge coupling unification constraints seem to imply that M_s cannot really be lower than 10 – 100 TeV.

An important question that arises in the context of unification via KK thresholds as well as its concrete realization via the TSSM is the issue of higher loop corrections. The point here is that even though the unified gauge coupling typically is small, the loop expansion parameter is still of order 1 for it is amplified by a large number of heavy KK modes running in the loops [12]. In the most general case it is therefore far from being obvious that the higher loop corrections are subleading. In fact, if one considers a generic KK compactification of a higher dimensional theory *without* supersymmetry, the higher loop corrections are as large as the one-loop threshold contribution, and therefore the very idea of gauge coupling unification via Kaluza-Klein thresholds in this context has no predictive power.

However, as was pointed out in [12], in the context of $\mathcal{N} = 1$ supersymmetric KK compactifications the situation is drastically different. In this paper we elaborate the discussions in [12] as well as [22] on the issue of higher loop corrections. In particular, we show that the higher loop corrections are subleading compared with the leading one-loop threshold contribution. Here we give two different ways of arriving at this result. The first argument is purely field theoretic and is along the lines of that given in [12], albeit the argument we give in this paper is somewhat simpler. The key observation underlying this argument is that the heavy KK modes (in certain orbifold compactifications we consider in this paper) have extended, namely, $\mathcal{N} = 2$ supersymmetry. This statement does not only concern the heavy KK spectrum, but also the *interactions* involving only heavy KK modes - the three- and four-point couplings of heavy KK states are $\mathcal{N} = 2$ supersymmetric. This then implies that the leading (in powers of the relevant number of the KK modes) contribution at a given higher loop order vanishes due to certain $\mathcal{N} = 2$ supersymmetric cancellations (recall that the gauge coupling is not renormalized beyond two loops in $\mathcal{N} = 2$ supersymmetric theories). The second argument we give in this paper is string theoretic and utilizes the power of string perturbation techniques. In fact, the string theoretic discussion we give in this paper allows one to arrive at a clear geometric interpretation of the above mentioned cancellations at higher loops via the D-brane picture as well as string world-sheet expansion in terms of Riemann surfaces of various topologies. In this sense, this argument is very much in the spirit of (but not exactly the same as) that employed in [26] to prove finiteness of certain large N gauge theories.

To illustrate the formal arguments mentioned above, we explicitly compute various two-loop corrections to the gauge couplings within the above framework. In fact, the cancellations of heavy KK contributions can be seen explicitly in these two-loop computations. We also discuss the general setup for performing higher loop computations in the KK theories of this type.

The rest of this paper is organized as follows. In section II we describe the setup for discussing higher loop corrections to the gauge coupling unification via Kaluza-Klein thresholds. In particular, we discuss certain orbifold compactifications of higher dimensional theories in this context. In section III we discuss one-loop unification via Kaluza-Klein thresholds. In particular, we give the general expression for the one-loop KK thresholds in the context of the orbifold theories discussed in section II. In section IV we give the field and string theoretic arguments which show that the underlying $\mathcal{N} = 2$ supersymmetry (at the heavy KK levels) indeed implies cancellation of the leading higher loop contributions. We also discuss the general setup for computing the subleading higher loop corrections. In section V we give explicit two loop computations in the above context. In section VI we remark on

various important issues in the context of unification via Kaluza-Klein thresholds.

II. SETUP

In this section we describe the setup for discussing higher loop corrections to the gauge coupling unification via Kaluza-Klein thresholds. In particular, we will discuss certain orbifold compactifications of higher dimensional theories in this context. For definiteness, we will focus on compactifications of six dimensional gauge theories. Generalization to other cases is completely straightforward.

Thus, consider a six dimensional gauge theory living in the world-volume of some set of coincident D5-branes⁹. Let the space transverse to the D5-branes be a K3 surface. Then we have $\mathcal{N} = 1$ supersymmetric gauge theory living in the world volume of the D5-branes. In the following discussion, for the most part, the directions transverse to the branes are not important. However, we will still keep track of them to have a clear geometric interpretation of the four dimensional gauge theory. Next, compactify two of the directions inside of the D5-branes on a two-torus T^2 . The low energy effective field theory is then given by the corresponding four dimensional $\mathcal{N} = 2$ supersymmetric gauge theory. Here by low energies we mean energies below the compactification scale. Thus, let $T^2 = S^1 \otimes S^1$, where both circles have the same radius R . Then the compactification scale is given by the mass scale of the first Kaluza-Klein threshold $1/R$. Below this scale we have an effective four dimensional gauge theory.

We would now like to obtain an $\mathcal{N} = 1$ supersymmetric gauge theory from this setup. To do this, we will mod out the above theory by a discrete orbifold group Γ . Here Γ must act on both T^2 and K3 such that $(T^2 \otimes \text{K3})/\Gamma$ is a Calabi-Yau three-fold with $SU(3)$ holonomy. Note that the action of Γ on T^2 must be crystallographic (or else it is not a symmetry of the theory). This restricts the allowed choices of Γ to Abelian cyclic groups $\mathbf{Z}_2, \mathbf{Z}_3, \mathbf{Z}_4, \mathbf{Z}_6$. In the following we will generally refer to Γ as $\mathbf{Z}_M$ with the understanding that M is restricted to the above values.

To discuss the spectrum of the above orbifold model, let us introduce some notations. First, let G be the gauge group of the original six dimensional gauge theory. Note that G can be a product group. Let the matter content of the six dimensional theory be given by hypermultiplets in various representations of G . We will collectively denote these matter hypermultiplets by $\tilde{\Phi}$. Each hypermultiplet can be written as $\tilde{\Phi} = \Phi \oplus \Phi'$, where Φ denotes a chiral multiplet (in the four dimensional sense) with left chirality, whereas Φ' denotes the corresponding anti-chiral multiplet with right chirality. Similarly, let $\tilde{V}$ be the gauge vector supermultiplet (of the four dimensional $\mathcal{N} = 2$ gauge theory) transforming in the adjoint of G . It can be written as $\tilde{V} = V \oplus \chi$, where V denotes the corresponding $\mathcal{N} = 1$ gauge vector supermultiplet (which contains a Weyl fermion with left chirality), whereas χ denotes the complement of V in $\tilde{V}$, that is, the $\mathcal{N} = 1$ anti-chiral supermultiplet (which contains a Weyl fermion with right chirality) transforming in the adjoint of G .

⁹Here we concentrate on D-brane theories, albeit our discussion is unmodified if we have some other type of branes.

Next, let us discuss the action of the orbifold group $\Gamma \approx \mathbf{Z}_M$ on various degrees of freedom. First, we will assume that the action of Γ on the gauge quantum numbers is trivial (that is we choose the corresponding gauge bundle to be trivial). Thus, orbifolding does *not* break the original gauge group G .

Now, let us discuss the breaking of the $\mathcal{N} = 2$ supersymmetry to $\mathcal{N} = 1$ by the orbifold group Γ . Let g be the generator of Γ . Then the action of g on the R-parity quantum numbers of states in $\tilde{V}$ and $\tilde{\Phi}$ is the following: $g|V\rangle = |V\rangle$, $g|\chi\rangle = \omega|\chi\rangle$, $g|\Phi\rangle = |\Phi\rangle$, $g|\Phi'\rangle = \omega^{-1}|\Phi'\rangle$, where $\omega \equiv \exp(2\pi i/M)$. Geometrically this can be understood from the fact that (up to an obvious choice of convention) g acts on the complex coordinate z_1 on T^2 as $gz_1 = \omega z_1$, while its action on the holomorphic two-form $\Omega_2 \equiv dz_2 \wedge dz_3$ on K3 is given by $g\Omega_2 = \omega^{-1}\Omega_2$. Here z_2, z_3 are the complex coordinates on K3. Note that the holomorphic three-form $\Omega_3 \equiv dz_1 \wedge dz_2 \wedge dz_3$ is invariant under the action of Γ , which is nothing but the condition that the quotient $(T^2 \otimes \text{K3})/\Gamma$ is a Calabi-Yau three-fold.

Finally, we turn to the action of Γ on the Kaluza-Klein modes corresponding to the compactification on T^2 . Let γ^{ij} , $i, j = 1, 2$, be the inverse metric¹⁰ on T^2 . Then the masses of the KK modes are given by:

$$M_{\mathbf{m}}^2 = \gamma^{ij} m_i m_j . \quad (1)$$

Here $\mathbf{m} \equiv (m_1, m_2)$, and m_1, m_2 are the integer KK momenta corresponding to the a -cycle respectively b -cycle on T^2 . The action of the generator g of Γ on the KK momenta is given by the corresponding $2\pi/M$ rotation: $g|\mathbf{m}\rangle = |\theta\mathbf{m}\rangle$. Here θ is the corresponding rotation matrix: $\theta : m_i \rightarrow \theta_i^j m_j$. Note that θ^M is an identity matrix.

The zero KK modes with $\mathbf{m} = (0, 0)$ are invariant under the action of g . At heavy KK levels with $\mathbf{m} \neq (0, 0)$ we can form linear combinations such that the action of g is diagonal. These are given by ($\ell = 0, 1, \dots, M-1$):

$$|\mathbf{m}; \ell\rangle \equiv \frac{1}{\sqrt{M}} \sum_{k=0}^{M-1} \omega^{-\ell k} |\theta^k \mathbf{m}\rangle . \quad (2)$$

Note that $g|\mathbf{m}; \ell\rangle = \omega^\ell |\mathbf{m}; \ell\rangle$.

Now we are ready to determine the spectrum of the orbifold theory. We must project onto the states invariant under the action of Γ , that is, onto the g -invariant states. These are given by ($\mathbf{m} \neq (0, 0)$)

$$\begin{aligned} & |(0, 0)\rangle \otimes |V\rangle , \quad |(0, 0)\rangle \otimes |\Phi\rangle , \\ & |\mathbf{m}; 0\rangle \otimes |V\rangle , \quad |\mathbf{m}; M-1\rangle \otimes |\chi\rangle , \\ & |\mathbf{m}; 0\rangle \otimes |\Phi\rangle , \quad |\mathbf{m}; 1\rangle \otimes |\Phi'\rangle . \end{aligned}$$

Here, to avoid overcounting, independent choices of $\mathbf{m}$ are restricted to the appropriate conjugacy classes each containing M elements such that they are all related by the corresponding θ rotations.

¹⁰Thus, in the case of a square torus $T^2 = S^1 \otimes S^1$ the inverse metric reads: $\gamma^{11} = 1/R_1^2$, $\gamma^{22} = 1/R_2^2$, and $\gamma^{12} = \gamma^{21} = 0$, where R_1 and R_2 are the radii of the two circles.

Note that the massless modes are $\mathcal{N} = 1$ supersymmetric. However, the massive KK modes form $\mathcal{N} = 2$ supermultiplets. Thus, for each choice of $\mathbf{m}$ (in the corresponding conjugacy class), $(|\mathbf{m}; 0\rangle \otimes |V\rangle) \oplus (|\mathbf{m}; M-1\rangle \otimes |\chi\rangle)$ forms a (short) massive $\mathcal{N} = 2$ supersymmetric vector multiplet. Similarly, $(|\mathbf{m}; 0\rangle \otimes |\Phi\rangle) \oplus (|\mathbf{m}; 1\rangle \otimes |\Phi'\rangle)$ forms a massive $\mathcal{N} = 2$ hypermultiplet. In fact, this statement is not just about the massive KK spectrum, but also holds for massive KK *interactions* as well. More concretely, it is not difficult to see that in the orbifold theory the three- and four-point couplings involving only heavy KK modes are $\mathcal{N} = 2$ supersymmetric with strengths rescaled by $1/\sqrt{M}$ respectively $1/M$ compared with the corresponding couplings in the parent $\mathcal{N} = 2$ theory. As we will see in the next section, this fact has important implications for higher loop corrections to the gauge coupling renormalization. In particular, it implies that the higher loop diagrams with only heavy KK modes running in the loops vanish due to $\mathcal{N} = 2$ supersymmetry. Thus, for a higher loop correction to be non-vanishing, it is required that it involves at least one massless line inside of the loops. As we have already mentioned, massless states do *not* have $\mathcal{N} = 2$ supersymmetry but are only $\mathcal{N} = 1$ supersymmetric so that the corresponding higher loop contributions are (generically) non-vanishing. Nonetheless, as we point out in the next section, such higher loop diagrams are subleading compared with the one-loop gauge coupling renormalization due to the KK thresholds. This is the key reason why the gauge coupling unification via KK thresholds is meaningful in the present setup.

Before we end this section, we would like to point out that we can introduce additional massless matter fields in the above picture such that they do not have heavy KK counterparts. We will collectively denote these fields by ϕ . Thus, the ϕ fields can be localized at the fixed points of the orbifold. That is, they are truly $3+1$ dimensional in contrast, say, to the fields $\tilde{V}$ and $\tilde{\Phi}$ which can be viewed as propagating in $5+1$ dimensions (two of which are compact). The fields ϕ then only have $\mathcal{N} = 1$ supersymmetry, and their couplings to other fields including heavy KK modes are also $\mathcal{N} = 1$ supersymmetric. One of the reasons for introducing such localized matter fields is that their presence is typically desirable in concrete phenomenological model building.

III. UNIFICATION

In this section we discuss one-loop threshold corrections to gauge coupling renormalization due to the heavy KK modes. In particular, we give the general expression for the KK thresholds in the context of the orbifold theories discussed in the previous section. We then apply these results to one-loop gauge coupling unification via Kaluza-Klein thresholds.

To begin with, let us clarify what we mean by gauge coupling unification. Suppose G is a product gauge group: $G = \otimes_a G_a$, where G_a are simple subgroups (which can be Abelian or non-Abelian). At the tree level the gauge couplings α_a for all subgroups are assumed to be the same: $\alpha_a \equiv \alpha$. In string theory this is the case if the entire gauge group comes from the same set of coincident branes. In fact, the tree-level “unified” gauge coupling α is given by (here we are using the conventions of [23]):

$$\alpha = g_s/2v_{p-3} . \quad (3)$$

Here g_s is the (ten dimensional) string coupling, and v_{p-3} measures the size of the $p-3$ compact spatial directions inside of the Dp -branes:

$$v_{p-3} \equiv V_{p-3}(M_s/2\pi)^{p-3} \equiv (RM_s)^{p-3} , \quad (4)$$

where V_{p-3} is the actual volume of these $p-3$ compact directions, and $M_s = 1/\sqrt{\alpha'}$ is the string scale. Note that here R is understood only as an “effective size” of compactification, and in general it need not coincide with the actual linear dimension(s) of the compactification space.

The above tree-level relations are subject to radiative corrections. Here we focus on one-loop corrections to the gauge couplings α_a . In particular, we are interested in gauge couplings at low energies $\mu \ll 1/R < M_s$. These gauge couplings $\alpha_a(\mu)$ depend upon the energy scale μ . More concretely, the energy scale dependent one-loop corrections to the gauge couplings come from the corresponding infra-red (IR) divergences due to the massless modes propagating in the loop. Then μ plays the role of the IR cut-off. This gives precisely the familiar field theoretic logarithmic evolution of gauge couplings. There are also energy scale *independent* corrections due to various *thresholds* corresponding to massive modes (with masses $\gg \mu$) propagating in the loop. These thresholds come from the heavy KK modes as well as string states such as string oscillator modes. At one loop we have:

$$\alpha_a^{-1}(\mu) = \alpha^{-1} + \frac{b_a}{2\pi} \ln \left(\frac{M_s}{\mu} \right) + \Delta_a , \quad (5)$$

where the one-loop β -function coefficients b_a correspond to the massless $\mathcal{N} = 1$ supersymmetric modes: $b_a = b_a(V) + b_a(\Phi) + b_a(\phi)$ (see section II for notations). Here we have chosen the ultra-violet (UV) cut-off to be the string scale M_s . This is equivalent to identifying the gauge coupling unification scale with the string scale. In particular, if we start from the low energy gauge couplings, we can run them up to the string scale by “integrating in” the heavy KK modes as we go to higher and higher energies. Then, with the above convention, the gauge couplings unify at the string scale M_s subject to the appropriate choice of the subtraction scheme. In the above approach, as we will see in a moment, the choice of the subtraction scheme affects the threshold corrections Δ_a .

In the following we are interested in the regime where $(RM_s)^{p-3} \gg 1$. In this case the threshold corrections due to the KK modes are large and dominate the string thresholds which generically are of order 1 (or smaller) [16]. We will therefore focus on the KK thresholds.

As we will see in a moment, the KK threshold computation in the $\mathcal{N} = 1$ supersymmetric orbifold theory reduces to that in the parent $\mathcal{N} = 2$ gauge theory. Let us therefore first consider the $\mathcal{N} = 2$ gauge theory with the superfields $\tilde{V}$ and $\tilde{\Phi}$ which arises upon compactifying the corresponding six dimensional $\mathcal{N} = 1$ supersymmetric gauge theory (living in the world-volume of D5-branes) on T^2 . In the following, for the sake of notational convenience, we will be a bit more general and treat it as a compactification of a $p+1$ dimensional theory (living in the world-volume of Dp-branes) on a $(p-3)$ -torus T^{p-3} . Note that p here can take values $p = 4, 5$, and in the former case we have a compactification of a five dimensional gauge theory on a circle (here the corresponding orbifold group can only be $\mathbf{Z}_2$). In these notations the KK spectrum of the $\mathcal{N} = 2$ theory consists of states with masses $M_{\mathbf{m}}^2 = \gamma^{ij} m_i m_j$ ($\mathbf{m} \equiv (m_1, \dots, m_{p-3})$, $m_i \in \mathbf{Z}$) with quantum numbers $\tilde{V}$ and $\tilde{\Phi}$. All of the states including the massless ones are $\mathcal{N} = 2$ supersymmetric, and only contribute to the renormalization of the low energy gauge couplings at one loop - perturbatively there are no corrections to

the gauge couplings beyond one loop, which is due to $\mathcal{N} = 2$ supersymmetry. Moreover, in the D-brane context *no other* states contribute to the gauge coupling renormalization. This follows from the fact that in perturbative open string theory only BPS states can renormalize gauge couplings [24,17]. In six dimensional $\mathcal{N} = 1$ open string theories the only BPS states are the massless modes, while all the other states are non-BPS as they come in six dimensional $\mathcal{N} = 2$ (that is, four dimensional $\mathcal{N} = 4$) supermultiplets. The latter, however, do not renormalize gauge couplings (in the four dimensional gauge theory arising upon compactification on T^2). Therefore, we are left only with the KK threshold contributions, which we are going to discuss next.

The computation of the gauge coupling renormalization in this $\mathcal{N} = 2$ gauge theory can be performed entirely within the field theory approach. In particular, we will treat this theory as a gauge theory with a UV cut-off Λ containing massless modes plus massive KK states. The role of the cut-off Λ is to restrict the heavy KK modes to a finite subset which essentially amounts to discarding those with masses larger than Λ . More precisely, one can employ the standard Coleman-Weinberg prescription which gives the following simple result [22]:

$$\alpha_a^{-1}(\mu) = \alpha_a^{-1}(\Lambda) + \frac{\tilde{b}_a}{4\pi} \int_{(\xi\Lambda)^{-2}}^{(\xi\mu)^{-2}} \frac{dt}{t} \sum_{\mathbf{m}} \exp(-\pi t M_{\mathbf{m}}^2) . \quad (6)$$

Here μ and Λ are the IR respectively UV cut-offs, and we have parametrized the subtraction scheme dependence by ξ . The one-loop β -function coefficients $\tilde{b}_a$ are those of the $\mathcal{N} = 2$ theory: $\tilde{b}_a = b_a(\tilde{V}) + b_a(\tilde{\Phi})$ (see section II for notations). Next, we will identify Λ with the string scale M_s , and the gauge couplings $\alpha_a(\Lambda)$ with the “unified” gauge coupling α . Note that this is the only place where string theory becomes relevant - as we will discuss in the next section, for the above prescription to be meaningful, we must assume that above the cut-off the theory is finite, which is precisely what we expect above the string scale where string theory description takes over. With these identifications, we will obtain the expected logarithmic evolution of the low energy ($\mu \ll 1/R < M_s$) gauge couplings $\alpha_a(\mu)$. This logarithmic contribution comes from the massless modes with $\mathbf{m} = 0$:

$$\alpha_a^{-1}(\mu) = \alpha^{-1} + \frac{\tilde{b}_a}{2\pi} \ln \left(\frac{M_s}{\mu} \right) + \tilde{\Delta}_a . \quad (7)$$

The IR *finite* threshold corrections $\tilde{\Delta}_a$ are due to the massive KK modes with $\mathbf{m} \neq (0, \dots, 0)$. The leading contribution (in the regime $(RM_s)^{p-3} \gg 1$) to $\tilde{\Delta}_a$ can be readily evaluated using the Poisson resummation, and the result is given by [22]:

$$\tilde{\Delta}_a = \frac{\tilde{b}_a}{2\pi} \frac{\xi^{p-3}}{p-3} (RM_s)^{p-3} - \frac{\tilde{b}_a}{2\pi} \ln(RM_s) + \mathcal{O}(1) . \quad (8)$$

Here for the sake of simplicity in the case of T^2 we assume that the metric γ_{ij} on T^2 satisfies $\gamma_{11} \sim \gamma_{22} \sim \det^{1/2}(\gamma_{ij}) = R^2$ (that is, the complex structure on T^2 is “of order 1”). Note that the subtraction scheme dependent parameter ξ cannot be determined within these considerations alone¹¹. However, in a given theory ξ affects the unification scale M_s (for

¹¹In [5] the choice of the subtraction scheme was such that $\xi^{p-3} = \pi^{(p-3)/2} / \Gamma((p-1)/2)$.

given values of the low energy gauge couplings), and it could, therefore, be determined by fixing M_s via some other low energy quantity (such as the Newton's constant G_N).

Now we are ready to determine the threshold corrections Δ_a in the $\mathcal{N} = 1$ orbifold theory. In fact, we simply have

$$\Delta_a = \tilde{\Delta}_a/M . \quad (9)$$

This follows from the fact that the number of the massive KK modes in the $\mathcal{N} = 1$ theory is M times smaller than in the parent $\mathcal{N} = 2$ theory, which is due to the $\mathbf{Z}_M$ orbifold projection. Note that, as we have already mentioned, in (9) we are ignoring other $\mathcal{O}(1)$ threshold corrections, namely, those due to heavy string modes.

IV. HIGHER LOOP CORRECTIONS

In the previous section we discussed one-loop renormalization of gauge couplings due to KK thresholds. In this section we address the issue of higher loop corrections. In particular, *a priori* it might seem that higher loops would destroy the one-loop prediction for the low energy gauge couplings obtained from the assumption that they unify at the string scale, or, equivalently, one-loop unification of gauge couplings via Kaluza-Klein thresholds. If so, then the entire framework would have no predictive power.

There are various ways of thinking about higher loop effects, some of them being more precise than others. Thus, one might naively argue that the theories discussed in section II are higher dimensional and therefore non-renormalizable. This would imply that one has no control over higher loop corrections. Fortunately, however, this naive argument is too naive. The key observation here is that the regimes we are dealing with are such that the theories under consideration never become higher dimensional. To make this statement more precise, let us recall the tree-level relation between the four dimensional gauge coupling α , the string coupling g_s , the string scale M_s (which is nothing but the UV cut-off in this context), and the compactification “size” R :

$$\alpha = g_s/2(RM_s)^{p-3} . \quad (10)$$

Note that we are working in the regime where the lowest KK threshold is below the string scale. In fact, it is much below the string scale: $(RM_s)^{p-3} \gg 1$. If we take R to be smaller than $1/M_s$, then the description in terms of D p -branes becomes inadequate - we have to “T-dualize” into a more adequate description in terms of lower dimensional branes. Thus, if all $p - 3$ extra spatial dimensions become smaller than $1/M_s$, the adequate description is in terms of D3-branes. Now, if we take R to be large compared with $1/M_s$, naively this might seem to be sufficient to go into the decompactification regime where the theory essentially becomes $p+1$ dimensional. However, we wish to keep the four dimensional gauge coupling α fixed, so for fixed g_s we would have to decrease M_s while increasing R such that $RM_s(\gg 1)$ is fixed. This implies that in the decompactification limit $R \rightarrow \infty$ the UV cut-off $\Lambda = M_s$ of the theory goes to zero. That is, the theory never becomes a higher dimensional theory (which would be non-renormalizable) with a finite cut-off. Another way of phrasing this statement is that we are dealing with four dimensional gauge theories with a large but *finite* number N of heavy KK modes. This number N is determined by the cut-off *vs.* the first

KK threshold ratio. That is, we can define N to be given by $N = (RM_s)^{p-3}$. Note that here we are implicitly making the assumption that above the UV cut-off scale M_s the theory is actually finite or else truncation of the KK modes to a finite subset could not be justified. This assumption, however, holds in the string theory context, so as long as we bear in mind that these theories are meaningful only if we embed them in a larger theory such as string theory, the above naive arguments about these theories being higher dimensional and non-renormalizable do not hold.

A priori there is, however, a more serious worry which can be made precise. Thus, consider a generic KK compactification (which, in particular, need not be supersymmetric). Naively it might seem that as long as the “unified” gauge coupling α is small the higher loop corrections are negligible. This is, however, not the case - the true loop expansion parameter is *not* $\alpha/2\pi$ (which would be the case if the effective field theory description in terms of just the light modes were adequate all the way up to the string scale). Rather, the correct expansion parameter is related to the string coupling g_s . This is due to the fact that although each KK mode (including the light modes) couples with the strength of order α , there are many, namely, N KK modes propagating inside of the loops. The true loop expansion parameter is therefore¹²

$$\lambda \equiv \alpha N / 2\pi . \quad (11)$$

This is analogous to considering the effective 't Hooft's coupling in large N gauge theories [25]. As pointed out in [12], in concrete phenomenological applications the effective coupling λ can be of order 1. If so, then generically one would expect that higher loop threshold corrections are just as large as one-loop thresholds, that is, that this framework lacks any predictive power.

However, in [12] it was pointed out that supersymmetry saves the day. That is, in the case of non-supersymmetric theories the above argument indeed shows that one has no control over higher loop effects. In the supersymmetric case, however, there are subtle cancellations at higher loops such that one loop thresholds are always dominant. Here we would like to reiterate the field theoretic argument of [12], and give another way of arriving at the same result using the power of string perturbation techniques. In the next section we will give explicit two-loop computations which illustrate the general arguments presented in the remainder of this section.

A. Field Theory Argument

In this subsection we first review (a simplified version of) the argument of [12] which shows that higher loop corrections to the gauge coupling unification via KK thresholds are

¹²This definition is consistent with the fact that in the closed string sector the loop expansion parameter is $(g_s/4\pi)^2 = \lambda^2$, while in the open string sector it is $n_D(g_s/4\pi) = n_D\lambda$. Here n_D is the number of Dp-branes. In the gauge theory discussion it arises through the corresponding β -function coefficients. In fact, the statement that $n_D\lambda$ is the loop expansion parameter in the open string sector is essentially precise. A more precise statement is that the L -th loop is weighted by $b_{L-1}\lambda^L$ just as is the case in the gauge theory language.

indeed subleading compared with the leading one-loop contribution. In the following for the sake of simplicity we will omit the subscript a labelling the corresponding gauge subgroups G_a . On general grounds we expect that the renormalized gauge coupling at scales $\mu \ll 1/R$ is given by

$$\alpha^{-1}(\mu) = \alpha^{-1} + f(\mu) + \Delta , \quad (12)$$

where Δ is the (μ -independent) contribution due to heavy KK thresholds, whereas $f(\mu)$ corresponds to the gauge coupling running. Here we are interested in estimating the sizes of both $f(\mu)$ and Δ . The L -loop order contribution $\alpha_L^{-1}(\mu)$ to $\alpha^{-1}(\mu)$ can be schematically written as

$$\alpha_L^{-1}(\mu) = \alpha^{-1} \sum_{m=0}^L c_{m,L} N^m \left(\frac{\alpha}{2\pi} \right)^L , \quad (13)$$

where m counts the number of the loop propagators corresponding to heavy KK modes with *independent* KK momentum summations. (Note that the KK momentum must be conserved inside of the loop diagram as the two external gauge boson lines carry zero KK momentum.) The enhancement factor N^m arises due to N heavy KK states propagating in each of the m internal lines. The coefficients $c_{m,L}$ generically depend on the energy scale μ via the appropriate IR cut-off. Note that for $N \gg 1$ the coefficients $c_{m,L} \sim 1$ (or smaller) for energy scales not too much smaller than $1/R$. (Thus, here and in the following we will treat logarithms of the type $\ln(RM_s)$ as being of order 1.)

For $\lambda \sim 1$, naively one might expect a large (that is, of order α^{-1}) contribution to $\alpha^{-1}(\mu)$ coming from the term $m = L$ in (13). (Note that this term is μ -independent. More precisely, it is finite in the limit $\mu \rightarrow 0$. This is because all of the internal propagators in this case correspond to heavy KK states.) However, for $L \geq 2$ this contribution actually vanishes due to $\mathcal{N} = 2$ supersymmetry of heavy KK modes. Indeed, as we have already pointed out in section II, the spectrum of heavy KK modes is $\mathcal{N} = 2$ supersymmetric. Moreover, their interactions (that is, those involving only heavy KK modes but no massless states) are also $\mathcal{N} = 2$ supersymmetric. This then implies that all diagrams with two external gauge bosons involving only heavy KK modes inside of the loops vanish for gauge couplings are not renormalized beyond one loop in $\mathcal{N} = 2$ theories. Thus, non-vanishing contributions can only come from diagrams involving at least one massless internal propagator (with massless modes corresponding to V, Φ, ϕ). These diagrams, however, are suppressed by additional powers of N . Thus, we have

$$\alpha_0^{-1} = \alpha^{-1} \sim N , \quad (14)$$

$$\alpha_1^{-1}(\mu) \sim N , \quad (15)$$

$$\alpha_{L>1}^{-1}(\mu) \sim \lambda^{L-1} \sim 1 , \quad (16)$$

where the estimates here should be understood symbolically (that is, we are suppressing the μ dependence in the corresponding contributions). This implies that higher loop contributions to the gauge coupling renormalization are subleading compared with the one-loop contribution, although they are of order 1 so we have to worry about the perturbative “convergence” issues. However, as was argued in [12] using properties of the holomorphic gauge coupling in $\mathcal{N} = 1$ supersymmetric theories, we expect the perturbation series to converge

for $\lambda < \eta_c$, where $\eta_c \sim 1$ is a model dependent convergence radius (which, in particular, depends on the superpotential). The resummed higher loop corrections are then still expected to be of order 1 and therefore subleading compared with the one-loop contribution [12].

In the cases where λ is somewhat smaller than 1 the perturbation theory can be trusted. In particular, in various phenomenologically applications one might wish to compute higher loop corrections to the gauge coupling unification. In section V we will setup the general framework for computing two-loop corrections and present various explicit calculations at the two-loop order. Here, however, we would like to point out a set of simple rules which seem to be useful in computing higher loop corrections in $\mathcal{N} = 1$ orbifold theories using the corresponding computations in the parent $\mathcal{N} = 2$ theories.

It is convenient to normalize gauge couplings so that they are the same in both the $\mathcal{N} = 1$ and $\mathcal{N} = 2$ theories. At the same time we will take the T^{p-3} in the case of the $\mathcal{N} = 2$ theory to be the same as that appearing in the orbifold T^{p-3}/Γ in the $\mathcal{N} = 1$ theory. Then we have $\text{Vol}(T^{p-3}/\mathbf{Z}_M) = V_{p-3}/M$, where V_{p-3} is the volume of T^{p-3} . To have the low energy gauge couplings the same in the two theories, we then must take $\tilde{g}_s = M g_s$, where $\tilde{g}_s$ and g_s are the string couplings in the $\mathcal{N} = 2$ respectively $\mathcal{N} = 1$ theories.

Having fixed the relative normalization for the gauge couplings in the two theories, let us now discuss three- and four-point couplings involving various states. As we have already pointed out in section II, the three-point couplings involving only heavy KK states are $1/\sqrt{M}$ times weaker in the $\mathcal{N} = 1$ theory compared with those in the parent $\mathcal{N} = 2$ theory. (It then follows that the corresponding factor for the four-point couplings involving only heavy KK states is $1/M$.) As to the three-point couplings involving some massless states with V, Φ quantum numbers, they are not affected by the orbifold projection. That is, if there is at least one massless state in a three-point coupling, then this coupling is the same as in the parent $\mathcal{N} = 2$ theory. On the other hand, the four-point couplings involving one massless state are reduced by $1/\sqrt{M}$, while the four-point couplings involving two massless states are unaffected. Using these rules we can obtain some useful information about the relation between various diagrams in the two theories. Thus, consider an L -loop diagram with $m > 0$ internal propagators corresponding to heavy KK modes with independent KK momentum summations. For definiteness let us focus on diagrams involving only three-point functions. (The discussion straightforwardly generalizes to diagrams involving four-point functions as well.) It is not difficult to see that the value of this diagram in the orbifold theory is reduced by the factor $(1/M)^{2m-1}$ compared with the same diagram in the parent $\mathcal{N} = 2$ theory. This suppression factor arises as follows. Note that the heavy KK propagators contribute the suppression factor $(1/M)^m$ compared with the corresponding diagram in the parent $\mathcal{N} = 2$ theory for the number of the heavy KK modes in the orbifold theory is M times smaller than in the $\mathcal{N} = 2$ theory. On the other hand, there are $2(m-1)$ three-point functions involving only heavy KK modes, and each of these three-point functions contributes the suppression factor of $1/\sqrt{M}$. Putting all of this together, we obtain the above total suppression factor of $(1/M)^{2m-1}$. Note that internal lines corresponding to massless states are not modified compared with the parent $\mathcal{N} = 2$ theory except for the fact that some of the original $\mathcal{N} = 2$ massless states are absent in the $\mathcal{N} = 1$ theory due to the orbifold projection. This implies that to compute a given diagram with, say, $m > 0$, we can take the corresponding diagram in the $\mathcal{N} = 2$ theory, discard the terms involving massless states projected out by the orbifold action, and divide the resulting expression by M^{2m-1} . This will give us the answer for the

corresponding $\mathcal{N} = 1$ computation. In the case of diagrams involving only massless internal propagators (that is, for $m = 0$), there is no suppression factor (unlike in the $m > 0$ case), and to obtain the answer starting from the corresponding $\mathcal{N} = 2$ computation, we only need to discard terms involving states projected out by the orbifold action.

The above observations are very useful for simplifying higher loop computations in the orbifold theory once the corresponding computations have been done in the parent $\mathcal{N} = 2$ theory. This will become clear in the next section where we present some explicit two-loop computations. We will see that these simplifications are quite substantial since certain diagrams (or, more precisely, combinations thereof) vanish *even* if they contain massless internal propagators. On the other hand, the diagrams involving “twisted” states cannot be obtained by a simple orbifold reduction of the $\mathcal{N} = 2$ theory, therefore they must be computed on a model-to-model basis.

B. String Theory Argument

In this subsection we give a simple string theory argument which allows one to arrive at the above result by utilizing the power of string perturbation techniques. Let us start from the parent $\mathcal{N} = 2$ supersymmetric theory. In string theory language the perturbative expansion is in terms of Riemann surfaces with various topologies. For the sake of simplicity let us focus on oriented Riemann surfaces with b boundaries and g handles. In particular, for the sake of simplicity we will ignore diagrams with cross-caps which are straightforward to incorporate in the following discussion (see the next subsection for some relevant comments). Each boundary corresponds to D-branes, whereas handles correspond to closed string loops. Note that each diagram is weighted by g_s^{2g-2+b} , where g_s is the string coupling.

Let us first consider the one-loop open string diagram with two boundaries and no handles. This is the annulus amplitude. In the open string loop channel it is weighted by $g_s^0 N$. The enhancement by a factor of N is due to heavy KK modes propagating in the loop. Alternatively, we can view this diagram in the closed string tree channel where a closed string is exchanged between two D-branes whose boundary states we will denote by $|\widetilde{D}\rangle_0$. Note that in the closed string tree channel there is no enhancement due to exchange of some large number of states - all such states are at least as heavy as $\sim M_s$ (the winding modes are even heavier with masses $\sim RM_s^2 \gg M_s$). This implies that the enhancement of the annulus amplitude by a factor of N in the closed string tree channel is due to the normalization of D-brane boundary states:

$$|\widetilde{D}\rangle_0 \sim N^{1/2} |\widetilde{D}'\rangle_0 . \quad (17)$$

Here the boundary state $|\widetilde{D}'\rangle_0$ is normalized so that ${}_0\langle\widetilde{D}'|\widetilde{D}'\rangle_0 \sim 1$. Note that this normalization factor can be derived directly by computing the annulus amplitude in the open string loop channel, and then performing the open-closed duality transformation (that is, the transformation $t \rightarrow 1/t$, with t the proper time on the annulus). Indeed, in the open string loop channel we have the KK momentum sum which in the closed string tree channel produces the volume factor V_{p-3} upon the appropriate Poisson resummation (recall that $N \sim V_{p-3} M_s^{p-3}$).

Let us now consider a diagram with b boundaries and no handles. Note that this diagram corresponds to $L = b - 1$ loops in the open string channel. In the closed string tree channel it

can be viewed as b D-branes connected by a closed string tree with $b - 2$ three-point vertices corresponding to closed string interactions. Each of these vertices is weighted by the factor $g_s/(M_s^{p-3}V_{p-3})^{1/2} \sim g_s/N^{1/2}$, where V_{p-3} is the volume of the $p - 3$ compact directions inside of Dp -branes. Here we are ignoring the volume of transverse directions (that is, of K3, or, more precisely, of $K3/\Gamma$, which we assume to be of order 1 in the string units). Thus, such a diagram is weighted by

$$N^{b/2}(g_s/N^{1/2})^{b-2} \sim \alpha^{L-1}N^L . \quad (18)$$

Here on the l.h.s. the first factor $N^{b/2}$ corresponds to b D-branes (or, more precisely, their boundary states $|\widetilde{D}\rangle_0$ scaling as $\sim N^{1/2}$), whereas the second factor $(g_s/N^{1/2})^{b-2}$ comes from the closed string interaction vertices. On the r.h.s. we have taken into account that $g_s/N \sim \alpha$, and $L = b - 1$. Thus, this contribution corresponds to the term $m = L$ in the expression (13) applied to the $\mathcal{N} = 2$ theory.

Diagrams with handles correspond to subleading terms with $m < L$ in the language of (13). Indeed, each handle is accompanied by two three-point interaction vertices leading to the suppression by $g_s^2/N \sim \alpha^2 N$. That is, diagrams with b boundaries and g handles are weighted by

$$N^{b/2}(g_s/N^{1/2})^{b-2+2g} \sim \alpha^{L-1}N^{L-g} , \quad (19)$$

where we have taken into account that $L = b + 2g - 1$. Thus, we see that $m = L - g$ in the language of (13). Note that the coefficients $c_{m,L}$ in the $\mathcal{N} = 2$ gauge theory vanish for $L > 1$. That is, the only non-vanishing diagram in this case is that with $b = 2, g = 0$.

Next, let us consider the orbifold theory with $\mathcal{N} = 1$ supersymmetry. Here we have M different types of D-branes whose corresponding boundary states we will denote by $|D\rangle_k$, $k = 0, 1, \dots, M - 1$. The *untwisted* boundary states $|D\rangle_0$ are coherent states made of (left-right symmetric) untwisted sector closed string states. They are related to the boundary states in the parent $\mathcal{N} = 2$ theory via

$$|D\rangle_0 = \frac{1}{\sqrt{M}}|\widetilde{D}\rangle_0 . \quad (20)$$

This can be, for instance, seen by noting that the annulus amplitude in the orbifold theory is obtained from that in the parent theory by inserting the projection operator (here g is the generator of $\mathbf{Z}_M$)

$$\frac{1}{M} \sum_{k=0}^{M-1} g^k \quad (21)$$

into the trace over the Hilbert space of open string states (in the loop channel). The contribution with $k = 0$ then corresponds to the closed string exchange between the untwisted boundary states in the closed string tree channel. This is, however, the same as in the parent theory up to the overall factor of $1/M$.

As to the contributions corresponding to $k \neq 0$, these in the closed string tree channel map to closed string exchanges between the corresponding *twisted* boundary states $|D\rangle_k$, $k \neq 0$. The latter are coherent states made of twisted sector closed string states localized at

the *fixed points* of the orbifold. Note that unlike the untwisted boundary states, the twisted boundary states are normalized so that there is no $N^{1/2}$ enhancement factor. That is,

$${}_0\langle D|D\rangle_0 \sim N , \quad (22)$$

while for the twisted boundary states ($k \neq 0$) we have

$${}_k\langle D|D\rangle_k \sim 1 . \quad (23)$$

This can be seen as follows. First, since the closed string states making up the twisted boundary states are localized at fixed points, these boundary states cannot possibly know about the volume of the compact space - they are fixed by the local properties of the orbifold in the vicinity of the fixed points. In particular, even if we take the volume of the orbifold to infinity (that is, if we take $V_{p-3} \rightarrow \infty$), the twisted boundary states remain the same. Equivalently, we can arrive at the same conclusion by noticing that in the open string loop channel a non-trivial twist g^k leaves only the origin of the KK momentum lattice invariant. This implies that the corresponding character does *not* contain a sum over the KK momentum states. In particular, the open-closed duality transformation (which amounts to taking $t \rightarrow 1/t$, where t is the proper time on the annulus) does *not* produce any volume factors (upon the corresponding Poisson resummation). Note that the untwisted characters do contain the KK momentum sum in the open string loop channel, so in the closed string tree channel we get the volume factor V_{p-3} (which eventually gives rise to the $N^{1/2}$ normalization of the untwisted boundary states).

To summarize, we have untwisted boundary states with the enhanced normalization factor, and also the twisted boundary states without such an enhancement factor. The former are the same as in the $\mathcal{N} = 2$ theory (up to the overall factor of $1/\sqrt{M}$). The latter are those that carry the information about supersymmetry breaking.

Now we are ready to discuss higher loop corrections in the $\mathcal{N} = 1$ theory. Let us start with diagrams without handles. Out of b boundaries we can have b_U untwisted boundaries plus b_T twisted boundaries. Note that if $b_T = 0$, then for $b > 2$ (that is, at two or higher loops) the corresponding contribution to the gauge coupling renormalization vanishes. Indeed, the corresponding diagram is the same as in the parent $\mathcal{N} = 2$ supersymmetric theory up to an overall factor given by the appropriate power of M .

Thus, non-vanishing diagrams for $b > 2$ are those with at least one twisted boundary state. Note that the closed string states propagating along the b_T closed string tubes connecting these boundary states to the interior of the closed string tree are the corresponding twisted closed string states. Three-point couplings between twisted closed string states are weighted by g_s with no additional volume suppression factors¹³. Three-point couplings of two (g^k and g^{M-k}) twisted sector states with untwisted sector states are weighted by $g_s/N^{1/2}$ (just as in the case of three untwisted sector states). It is then not difficult to see that the diagrams involving twisted boundary states can at most be weighted by $g_s^{b-2} \sim \alpha^{L-1} N^{L-1}$.

¹³More precisely, this is the case for twisted states localized at the same fixed point. Couplings of states localized at different fixed points are exponentially suppressed with the volume. Such couplings, however, are not relevant for our discussion here.

For instance, a diagram with $b_U = 0$ with *all* (including those in the interior of the diagram) closed string tubes corresponding to twisted sector states would have such a weight. Other diagrams either have the same weight or are more suppressed (by additional factors of $1/N$). Here one must take into account that the “total twist” associated with a given diagram must be trivial, that is, if k_i , $i = 1, \dots, b$, label twists corresponding to the boundary states, we must have

$$\sum_{i=1}^b k_i = 0 \pmod{M} . \quad (24)$$

Thus, the diagrams that “know” about reduction of supersymmetry from $\mathcal{N} = 2$ to $\mathcal{N} = 1$ are all subleading (as they are weighted by $\alpha^{L-1} N^{L-1} \sim \lambda^{L-1} < 1$) compared with the leading one-loop diagram (with $b = 2$) weighted by $N \gg 1$. The reader can easily verify that adding handles does not change this conclusion. Thus, we have arrived at the result that the one-loop gauge coupling renormalization is dominant ($\sim N$) compared with the higher loop corrections (which are at best of order 1).

Here we would like to point out that the above argument is very much in the spirit of (but not exactly the same as) that employed in [26] to prove finiteness of certain large N gauge theories.

C. Comments

Before we end this section, we would like to make a few clarifying remarks. First, the string theory argument we gave in the previous subsection makes use of the by now well appreciated fact that string perturbation theory is a very efficient way of organizing various gauge theory diagrams. So string theory in this argument is only a tool - at the end of the day we must take a limit $M_s \rightarrow \infty$ (more precisely, $M_s \gg 1/R \gg \mu$), and the heavy string modes do not play any role except for providing a proper UV cut-off for the resulting gauge theory computation.

Next, in the argument of the previous subsection we only considered Riemann surfaces with boundaries and handles but no cross-caps. Cross-caps are boundary states corresponding to the orientifold planes. In the context of *perturbative* compactifications of open plus closed string theories the latter are required to cancel various tadpoles. Tadpole cancellation implies UV finiteness of the theory which, in the field theory language, would result in cancellation of all power-like threshold corrections to the gauge couplings. This is due to a cancellation between the annulus (a cylinder with two boundaries) and the Möbius strip (a cylinder with one boundary and one cross-cap) amplitudes [17,27]. In the closed string tree-channel this corresponds to cancellation of the tadpoles due to massless states corresponding to the zero winding modes that couple to D-branes wrapped on a torus.

Thus, as was already pointed out in [12,28], unification via Kaluza-Klein thresholds should be considered in the context of *non-perturbative* orientifolds where perturbative tadpoles do *not* cancel. The finiteness of the theory is then due additional *non-perturbative* states (which can arise at both massless as well as massive levels) which provide an effective cut-off in the theory just as in perturbative heterotic compactifications where the modular integration is restricted to the fundamental domain which excludes extreme UV momenta

(that is, those above a certain energy scale $\sim M_s$). In fact, (some) non-perturbative orientifolds, examples of which have been recently discussed in [29], can be viewed as hybrid compactifications sharing the features of both perturbative heterotic and orientifold compactifications albeit such compactifications are non-perturbative from both heterotic and orientifold viewpoints. As to the gauge theory computation, in this context it must be regularized by introducing a UV cut-off which we identify with the string scale. Such regulator (which does not arise in perturbative orientifolds [27]) is expected to arise in *non-perturbative* orientifolds once we take into account non-perturbative (from the orientifold viewpoint) string states which do not have a perturbative description in terms of, say, open strings ending on D-branes. These states are expected from Type I-heterotic duality - thus, already in ten dimensions there are infinitely many BPS states (charged under $\text{Spin}(32)/\mathbf{Z}_2$) on the heterotic side which cannot be seen perturbatively on the Type I side.

Thus, in the above setup, that is, in the context of non-perturbative orientifolds, gauge coupling renormalization can be computed in the gauge theory context assuming the appropriate cut-off at the string scale M_s (more precisely, at some scale of order M_s - we have to deal with the subtraction scheme ambiguities). The reason why this UV cut-off scale should be around M_s is clear - quantum gravity becomes important at energies of order of the (fundamental) Planck scale $M_P \sim M_s/g_s$, which is of order of the string scale for $g_s \sim 1$.

V. EXPLICIT TWO-LOOP COMPUTATIONS

In this section we present a computation of two-loop corrections to $U(1)$ gauge couplings, for the D5-brane orbifold models described in section II. This Abelian two-loop case is completely sufficient to illustrate some general features of higher loop corrections.

We will use the standard background field method and evaluate the corresponding photon vacuum polarization diagrams. Our starting point are the diagrams of $\mathcal{N} = 1$ supersymmetric QED in $D = 6$ dimensions. The diagrams describing the compactified $\mathcal{N} = 2$ parent theory contain internal propagators with momenta of the form $(p, \mathbf{m}/R)$, where $\mathbf{m}/R$ is the KK component, a two-dimensional lattice vector. The d^4p momentum loop integrals are UV divergent and require a cut-off. We will apply dimensional regularization and continue the divergent integrals to $d = 4 - \varepsilon$ dimensions. The UV divergences appear then as $1/\varepsilon$ poles; these are subtracted in the renormalization procedure, leaving a finite result which depends on the dimensional regularization scale Λ . The scale Λ is a UV cut-off which (in an appropriate subtraction scheme) can be identified with the string scale [16,30]. However, we are not interested here in the details of UV pole subtractions, therefore we consider “bare” diagrams without the UV counterterms.

The $\mathcal{N} = 2$ supersymmetric two-loop vacuum polarization diagrams (note that here $q^2 = \mu^2$)

$$\Pi_{\mu\nu}(q) = (q_\mu q_\nu - g_{\mu\nu} q^2) \Pi_2(q^2) \quad (25)$$

fall into 3 classes, F_2 , S_2 and Y_2 , involving respectively:

- hypermultiplet fermion loops;
- hypermultiplet scalar loops;
- SUSY Yukawa couplings of gauginos to hypermultiplets.

These diagrams are depicted in Fig.1. Note that all traces of gamma matrices and contractions of Lorentz indices are performed in $D = 6$ dimensions, ensuring the right counting of fermionic degrees of freedom as well as incorporating the Yukawa couplings of four-dimensional scalar photons to hypermultiplet fermions on the same footing as other gauge couplings. As a result of a straightforward computation¹⁴ we obtain the corresponding contributions to $\Pi_2(q^2)$ as the integrals over 3 Feynman parameters, x , y and z :

$$\begin{aligned} F_2 &= 0 , \\ S_2 &= \alpha\Gamma(\varepsilon) \frac{(4\pi N)^\varepsilon}{(2\pi)^3} \sum_{\mathbf{m}, \mathbf{n}} \int_0^1 \int_0^1 \int_0^1 dx dy dz \frac{2z(1-2zy)(1-z)^{-1+\varepsilon/2}[x(1-x)]^{-\varepsilon/2}}{\left[\frac{(\mathbf{m}+\mathbf{n}x)^2}{x(1-x)}(1-z) + \mathbf{n}^2 + q^2 R^2 zy(1-zy)\right]^\varepsilon} , \\ Y_2 &= -S_2 , \end{aligned} \quad (26)$$

where $N = (\Lambda R)^2$. Note that the diagrams F_2 are the same as the diagrams describing a version of $D = 4$ non-supersymmetric QED obtained by a toroidal compactification of the “minimal” $D = 6$ QED with one charged “electron”. These diagrams cancel among themselves hence there is no two-loop contribution to the β -function in such a theory. Furthermore the electron mass is not renormalized at one loop.

The sum of all three contributions,

$$\Pi_2(q^2) = F_2 + S_2 + Y_2 = 0 . \quad (27)$$

In this way, we find a complete cancellation of two-loop corrections to the gauge coupling. Of course, the cancellation of $1/\varepsilon$ poles is a consequence of the known result, $b_2 = 0$ in $\mathcal{N} = 2$ SUSY gauge theories. Our two-loop computation provides further evidence for the *complete* cancellation of higher loop corrections to gauge couplings.

We now proceed to the $\mathcal{N} = 1$ supersymmetric orbifold theory. As argued in the previous section, the leading large N contribution [$m = L = 2$ in the notation of (13)], which is due to diagrams involving *massive* KK excitations propagating in *all* internal lines, is absent. This is due to $\mathcal{N} = 2$ supersymmetry of the massive spectrum which is responsible for the cancellations encountered above. The subleading contribution, $m = 1$, arises from the diagrams involving one massless propagator while the $m = 0$ contribution is due to purely massless diagrams. However, here $\mathcal{N} = 2$ cancellations also result in the vanishing of certain contributions.

In the case under consideration, the two-loop diagrams involving *one* massless propagator with the Φ quantum numbers add up to zero. This can be seen as follows. First consider the corresponding diagrams in the $\mathcal{N} = 2$ theory where the massless states carry the $\tilde{\Phi}$ quantum numbers. If we ignore the external lines (corresponding to gauge bosons), then we have two more internal lines (both of which correspond to heavy KK modes) - one with $\tilde{\Phi}$ quantum numbers, and the other one with the $\tilde{V}$ quantum numbers. It is convenient to denote such a diagram by $\mathcal{A}(\tilde{\Phi}_0, \tilde{\Phi}_{\mathbf{m}}, \tilde{V}_{-\mathbf{m}})$, where subscripts indicate the KK momenta. In the $\mathcal{N} = 1$ language we have $\tilde{\Phi} = \Phi \oplus \Phi'$, and $\tilde{V} = V \oplus \chi$ (see section II), so that the above $\mathcal{N} = 2$ diagram can be written as:

¹⁴We used the symbolic manipulation program FORM Version 1.1 written by J. Vermaseren.

$$\begin{aligned}\mathcal{A}(\tilde{\Phi}_0, \tilde{\Phi}_{\mathbf{m}}, \tilde{V}_{-\mathbf{m}}) &= \mathcal{A}(\Phi_0, \Phi_{\mathbf{m}}, V_{-\mathbf{m}}) + \mathcal{A}(\Phi'_0, \Phi'_{\mathbf{m}}, V_{-\mathbf{m}}) + \\ &\quad \mathcal{A}(\Phi_0, \Phi'_{\mathbf{m}}, \chi_{-\mathbf{m}}) + \mathcal{A}(\Phi'_0, \Phi_{\mathbf{m}}, \chi_{-\mathbf{m}}) .\end{aligned}\tag{28}$$

Note that by the left-right symmetry ($\Phi \leftrightarrow \Phi'$) of the theory, we have $\mathcal{A}(\Phi_0, \Phi_{\mathbf{m}}, V_{-\mathbf{m}}) = \mathcal{A}(\Phi'_0, \Phi'_{\mathbf{m}}, V_{-\mathbf{m}})$, and $\mathcal{A}(\Phi_0, \Phi'_{\mathbf{m}}, \chi_{-\mathbf{m}}) = \mathcal{A}(\Phi'_0, \Phi_{\mathbf{m}}, \chi_{-\mathbf{m}})$. On the other hand, the above $\mathcal{N} = 2$ diagram vanishes: $\mathcal{A}(\tilde{\Phi}_0, \tilde{\Phi}_{\mathbf{m}}, \tilde{V}_{-\mathbf{m}}) = 0$. This implies that

$$\mathcal{A}(\Phi_0, \Phi_{\mathbf{m}}, V_{-\mathbf{m}}) = -\mathcal{A}(\Phi_0, \Phi'_{\mathbf{m}}, \chi_{-\mathbf{m}}) .\tag{29}$$

Next, let us see what happens to the above diagrams upon the $\mathbf{Z}_M$ orbifold projection (that is, let us discuss the corresponding diagrams in the context of the $\mathcal{N} = 1$ theory). We must discard the diagrams involving Φ'_0 as the latter states are projected out. We therefore conclude that after orbifolding the diagram $\mathcal{A}(\tilde{\Phi}_0, \tilde{\Phi}_{\mathbf{m}}, \tilde{V}_{-\mathbf{m}})$ reduces to

$$\mathcal{B} = \frac{1}{M} (\mathcal{A}(\Phi_0, \Phi_{\mathbf{m}}, V_{-\mathbf{m}}) + \mathcal{A}(\Phi_0, \Phi'_{\mathbf{m}}, \chi_{-\mathbf{m}})) = 0 .\tag{30}$$

Here the factor of $1/M$ arises due to the fact that in the $\mathcal{N} = 1$ theory we must restrict the summation over the KK modes with KK momenta in the corresponding conjugacy classes (see section II).¹⁵ On the other hand, the $\mathcal{N} = 1$ diagram $\mathcal{B}$ still vanishes due to (29). Thus, we have shown that the two-loop diagrams with only one massless internal line involving the states with the Φ quantum numbers indeed add up to zero.

The above discussion implies that non-vanishing diagrams with only one massless internal line must be such that the corresponding massless states carry the V quantum numbers. In the following, we will evaluate these diagrams; they involve one massless internal line with the V quantum numbers plus heavy KK lines with Φ, Φ' quantum numbers. The computation can actually be simplified by relating the corresponding diagrams in the $\mathcal{N} = 1$ theory to their $\mathcal{N} = 2$ counterparts. Let us therefore consider these $\mathcal{N} = 2$ diagrams which we will refer to as $\mathcal{A}(\tilde{\Phi}_{\mathbf{m}}, \tilde{\Phi}_{-\mathbf{m}}, \tilde{V}_0)$. In the $\mathcal{N} = 1$ language this $\mathcal{N} = 2$ diagram can be written as:

$$\begin{aligned}\mathcal{A}(\tilde{\Phi}_{\mathbf{m}}, \tilde{\Phi}_{-\mathbf{m}}, \tilde{V}_0) &= \mathcal{A}(\Phi_{\mathbf{m}}, \Phi_{-\mathbf{m}}, V_0) + \mathcal{A}(\Phi'_{\mathbf{m}}, \Phi'_{-\mathbf{m}}, V_0) + \\ &\quad \mathcal{A}(\Phi_{\mathbf{m}}, \Phi'_{-\mathbf{m}}, \chi_0) + \mathcal{A}(\Phi'_{\mathbf{m}}, \Phi_{-\mathbf{m}}, \chi_0) .\end{aligned}\tag{31}$$

Note that by the $\Phi \leftrightarrow \Phi'$ symmetry we have $\mathcal{A}(\Phi_{\mathbf{m}}, \Phi_{-\mathbf{m}}, V_0) = \mathcal{A}(\Phi'_{\mathbf{m}}, \Phi'_{-\mathbf{m}}, V_0)$, and $\mathcal{A}(\Phi_{\mathbf{m}}, \Phi'_{-\mathbf{m}}, \chi_0) = \mathcal{A}(\Phi'_{\mathbf{m}}, \Phi_{-\mathbf{m}}, \chi_0)$. On the other hand, the above $\mathcal{N} = 2$ diagram vanishes: $\mathcal{A}(\tilde{\Phi}_{\mathbf{m}}, \tilde{\Phi}_{-\mathbf{m}}, \tilde{V}_0) = 0$. This implies that

$$\mathcal{A}(\Phi_{\mathbf{m}}, \Phi_{-\mathbf{m}}, V_0) = -\mathcal{A}(\Phi_{\mathbf{m}}, \Phi'_{-\mathbf{m}}, \chi_0) .\tag{32}$$

Let us now extract the corresponding $\mathcal{N} = 1$ diagrams, which we will refer to as $\mathcal{C}$, surviving the orbifold projection. Since χ_0 is projected out by the $\mathbf{Z}_M$ orbifold, we have

$$\mathcal{C} = \frac{1}{M} (\mathcal{A}(\Phi_{\mathbf{m}}, \Phi_{-\mathbf{m}}, V_0) + \mathcal{A}(\Phi'_{\mathbf{m}}, \Phi'_{-\mathbf{m}}, V_0)) = \frac{2}{M} \mathcal{A}(\Phi_{\mathbf{m}}, \Phi_{-\mathbf{m}}, V_0) .\tag{33}$$

¹⁵This is precisely the factor $(1/M)^{2m-1}$ discussed in the previous section, specified to the case of $m = 1$.

Thus, we see that non-vanishing diagrams of this type are the same as those in $\mathcal{N} = 1$ supersymmetric QED (with massive charged superfields). Note that the corresponding diagram with all massless internal lines is given by

$$\mathcal{C}_0 = \mathcal{A}(\Phi_0, \Phi_0, V_0) . \quad (34)$$

Also, in practice, we found it most convenient to start from the original $\mathcal{N} = 2$ diagrams and subtract the contributions involving χ_0 component of the vector multiplet, *i.e.*, the scalar photon and one of the two photinos. Thus, in particular, we eliminated the contributions of scalar photons in S_2 and F_2 by subtracting the corresponding polarization components of the $D = 6$ gauge boson, and adjusted the overall factor in Y_2 to eliminate the redundant photino propagator. First, we write down the final result for purely massless diagrams, with all internal lines corresponding to $\mathbf{m} = \mathbf{n} = \mathbf{0}$:

$$\Pi^{\mathbf{0}}(q^2) = \alpha \frac{(\Lambda^2/q^2)^\varepsilon}{2(2\pi)^3} \left[\frac{1}{\varepsilon} + \ln(4\pi) - \gamma + \frac{7}{2} \right] , \quad (35)$$

where γ is the Euler's constant. It is easy to check that the coefficient of the $1/\varepsilon$ pole does correctly reproduce the two-loop coefficient of $\mathcal{N} = 1$ supersymmetric QED. The diagrams involving only one massless propagator are given by more complicated expressions involving Feynman parameter integrals such as in (26). However in the most interesting “low energy, large radius” limit, $\Lambda^2/q^2 \gg N \gg 1$, these integrals simplify and the result can be written as a momentum-independent correction:

$$\Pi^{\mathbf{m}} = \alpha \frac{2}{M} \sum_{\mathbf{m}} \frac{(N/\mathbf{m}^2)^\varepsilon}{2(2\pi)^3} \left[\frac{1}{\varepsilon} + \ln(4\pi) - \gamma + \frac{1}{2} \right] + \delta^{\mathbf{m}} . \quad (36)$$

In the above expression we have isolated the contribution

$$\delta^{\mathbf{m}} = -\alpha \frac{4}{M} \sum_{\mathbf{m}} \frac{(N/\mathbf{m}^2)^\varepsilon}{2(2\pi)^3} \left[\frac{1}{\varepsilon} + \ln(4\pi) - \gamma + \frac{1}{2} \right] , \quad (37)$$

which arises from the diagrams involving self-energy corrections to internal lines, and contains the UV divergence associated with the mass renormalization at the one-loop level. The finite part of $\delta^{\mathbf{m}}$ should be combined with the one-loop threshold correction. It renormalizes the mass scale of the KK excitations, *i.e.*, the compactification radius R , or, equivalently, the number N of the KK excitations below the UV cut-off scale. On the other hand, the first term in (36) represents a genuine two-loop threshold correction proportional to the two-loop β -function of $\mathcal{N} = 1$ supersymmetric theory describing $\mathcal{N} = 2$ KK excitations coupled to the $\mathcal{N} = 1$ massless gauge sector. Note that in the above two-loop expression for the heavy KK threshold contributions to the gauge coupling renormalization the summation over the KK momenta $\mathbf{m}$ is cut-off from above by $N = (\Lambda R)^2$. After the summation the leading two-loop threshold is of order $\alpha N \sim 1$, which is subleading compared with the leading one-loop threshold correction $\sim N$.

VI. DISCUSSIONS

In this section we would like to discuss various issues which, we believe, have important implications in phenomenological applications of gauge coupling unification via Kaluza-Klein

thresholds. Some of these points have been previously stressed in [12], while some others are new and concern various developments subsequent to [12].

One of the key points we would like to emphasize is that supersymmetry is crucial for the arguments of section IV to go through. In particular, without supersymmetry, namely, without $\mathcal{N} = 2$ supersymmetry at the heavy KK levels, the cancellations leading to the fact that higher loop corrections to the gauge couplings are suppressed compared with the leading one-loop threshold contribution would not be possible. That is, in the context of KK compactifications of higher dimensional theories without supersymmetry the very idea of gauge coupling unification via Kaluza-Klein thresholds would have no predictive power - higher loop corrections in such a setup would be as large as one-loop thresholds, hence no control over the former.

Even in the supersymmetric case the fact that higher loop corrections are subleading compared with the one-loop thresholds is only specific to gauge coupling renormalization. Thus, the same does *not* hold for higher point couplings. In particular, the Yukawa couplings receive large higher loop corrections, and therefore one no longer has predictions for Yukawa coupling unification such as the $b - \tau$ unification.

Since supersymmetry at the heavy KK levels is so crucial, one has to worry about the effects of supersymmetry breaking which must eventually take place. Thus, the supersymmetry breaking scale (by which we mean the typical scale of soft scalar masses) must be low enough compared with the KK threshold scale $1/R$ or else the corrections due to supersymmetry breaking at the heavy KK levels would be too large and spoil the above mentioned cancellations due to $\mathcal{N} = 2$ supersymmetry. This implies that the string scale $M_s \gg 1/R$ cannot really be brought down to a few TeV in this context. (In the particular model of [12] the relation between the string scale M_s and the KK threshold $1/R$ is fixed by the unification constraint to be $M_s \sim 40/R$ in the case of D4-branes, and $M_s \sim 6/R$ in the case of D5-branes.) In fact, these considerations suggest that the string scale cannot be lower than 10 – 100 TeV.

To implement gauge coupling unification via Kaluza-Klein thresholds in the phenomenological context a concrete model is required. Such a model, which in [28] was called TSSM (TeV-scale Supersymmetric Standard Model), was proposed in [12]. In fact, it was stressed in [12] that this model was the only solution to the unification constraints found in [12]. Here we would like to discuss some issues concerning this point.

First, in the MSSM the gauge coupling running is given by:

$$\alpha_a^{-1}(\mu) = \alpha_{GUT} + \frac{b_a^*}{2\pi} \ln \left(\frac{M_{GUT}}{\mu} \right) , \quad (38)$$

where $\alpha_{GUT} \approx 1/24$ is the unified gauge coupling, $M_{GUT} \approx 2 \times 10^{16}$ GeV [20] is the unification scale, and b_a^* , $a = 1, 2, 3$, are the $SU(3)_c \otimes SU(2)_w \otimes U(1)_Y$ β -function coefficients ($b_1^* = 33/5$, $b_2^* = 1$, $b_3^* = -3$, where we have used the standard normalization $\alpha_1 = (5/3)\alpha_Y$). In order for the (one-loop) unification via the KK thresholds to be just as precise as in the MSSM, it is required that the $\mathcal{N} = 2$ β -function coefficients $\tilde{b}_a$ at the massive KK levels (see section III) satisfy the following relation:

$$\nu_{ab} \text{ is independent of } a, b, \text{ where } \nu_{ab} \equiv \frac{\tilde{b}_a - \tilde{b}_b}{b_a^* - b_b^*} \text{ for } a \neq b. \quad (39)$$

This condition is satisfied in the TSSM. In fact, in the TSSM we have $\nu_{ab} = 1 \ \forall a \neq b$. As was explained in [12], the TSSM was the only solution found there for the unification constraint (39). Here we point out that the key assumption here is that all three gauge subgroups of $SU(3)_c \otimes SU(2)_w \otimes U(1)_Y$ come from the *same* set of coincident branes. In this case in the string theory context we expect all three gauge couplings to be the same at the string scale (which is identified with the unification scale).

Recently, in [31,32] a set of models different from the TSSM was discussed. According to [31,32], in these models the (one-loop) gauge coupling unification is as precise as in the MSSM (and, therefore, TSSM). Here, however, we would like to point out that there is *no* unifications in the models of [31,32]. The point here is that in [31,32] additional “solutions” to the unification constraints were found by relaxing the requirement that all the gauge subgroups come from the same set of coincident branes. For instance, *a priori* one could imagine that $SU(3)_c \otimes U(1)_Y$ subgroups are localized on the same set of coincident Dp -branes (with $p = 4, 5$), while $SU(2)_w$ is localized on a fixed point of the orbifold (such localization, for instance, could be achieved by having other branes stuck at the orbifold fixed points in the corresponding compact directions). However, in such cases the gauge couplings of $SU(3)_c \otimes U(1)_Y$ on the one hand, and $SU(2)_w$ on the other hand generically are *not* expected to be the same at the string scale. In certain cases equality of these gauge couplings can be achieved by some fine tuning, but there is *no* unification prediction here. In this sense such a scenario is no different from that proposed in [8] where various subgroups of the Standard Model gauge group come from different sets of branes (and then unification requires fine tuning of the corresponding compactification volumes). Thus, the TSSM of [12] (along with its straightforward variations) is indeed the only known solution that satisfied the unification constraints.

Here another remark is in order. Note that the consistent orbifold reduction of a higher dimensional gauge theory requires that the heavy KK spectrum is $\mathcal{N} = 2$ supersymmetric. Moreover, suppose we have chiral multiplets Φ at the massless level. Then at the massive level we must have the corresponding hypermultiplet (we have been referring to these hypermultiplets as $\tilde{\Phi}$) for *each* massless chiral multiplet (with certain degeneracy depending on the order M of the orbifold group $\mathbf{Z}_M$ and the choice of T^{p-3}). This requirement, which is necessary for the consistency (in particular, unitarity) of the theory at the heavy KK levels is *not* met by the models of [5]. Thus, these models are not completely consistent. Similarly, albeit the unification is not a property in models of [31,32], some care is needed when considering such constructions. In particular, if, say, $SU(2)_w$ is localized on a fixed point of the orbifold, so must be *all* the matter fields charged under $SU(2)_w$ (which is *not* the case in some of the models of [31]) - this is the consequence of the corresponding flux conservation requirement.

Finally, we would like to comment on the “UV sensitivity” issue in the context of gauge coupling unification via Kaluza-Klein thresholds. Thus, naively it might seem that a small shift in the low energy gauge couplings would spoil the unification prediction (whose sensitivity to such shifts is power-like), and therefore the gauge coupling unification is too “UV sensitive” to be predictive. However, such a viewpoint might be a bit misleading. Thus, consider the TSSM. If the unification occurs in the MSSM, then it also occurs in the TSSM, and just as precisely (at one loop) as in the MSSM. Higher loop corrections, as we have shown in this paper, are small. This implies that such a unification scenario is predictive in

the context of a concrete model, namely, the TSSM. This is essentially due to the fact that the TSSM *explains* [12] why the usual logarithmic unification in the MSSM is not an accident (assuming that the TSSM is the correct description of physics above the electroweak scale), but is rather related to the lack of experimental data which prompts one to assume the standard “desert scenario” and extrapolate the gauge couplings all the way to the GUT scale M_{GUT} .

ACKNOWLEDGMENTS

We would like to thank Costas Bachas, Gia Dvali and Pran Nath for valuable discussions. This work was supported in part by the grant NSF PHY-96-02074. The work of Z.K. was also partially supported by the DOE 1994 OJI award. Z.K. would also like to thank Albert and Ribena Yu for financial support.

FIGURES




FIG. 1. Various two-loop gauge theory diagrams discussed in section V. Solid lines are hypermultiplet scalars, dashed lines are hypermultiplet fermions, double-dashed lines are gauginos, and wavy lines are gauge bosons. Note that tadpole-like contributions to the scalar propagators are not shown. Also, one should weight the above diagrams with the appropriate symmetry factors.

REFERENCES

- [1] J. Polchinski, Phys. Rev. Lett. **75** (1995) 4724.
- [2] E. Witten, Nucl. Phys. **B471** (1996) 135.
- [3] J. Lykken, Phys. Rev. **D54** (1996) 3693.
- [4] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. **B429** (1998) 263.
- [5] K.R. Dienes, E. Dudas and T. Gherghetta, Phys. Lett. **B436** (1998) 55; Nucl. Phys. **B537** (1999) 47; hep-ph/9807522.
- [6] Z. Kakushadze, Phys. Lett. **B434** (1998) 269; Nucl. Phys. **B535** (1998) 311; Phys. Rev. **D58** (1998) 101901.
- [7] I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. **B436** (1998) 257.
- [8] G. Shiu and S.-H.H. Tye, Phys. Rev. **D58** (1998) 106007.
- [9] Z. Kakushadze and S.-H.H. Tye, Phys. Rev. **D58** (1998) 126001.
- [10] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, hep-ph/9807344.
- [11] Z. Kakushadze and S.-H.H. Tye, Nucl. Phys. **B548** (1999) 180.
- [12] Z. Kakushadze, Nucl. Phys. **B548** (1999) 205.
- [13] I. Antoniadis, Phys. Lett. **B246** (1990) 377.
- [14] V. Rubakov and M. Shaposhnikov, Phys. Lett. **B125** (1983) 136;
A. Barnaveli and O. Kancheli, Sov. J. Nucl. Phys. **52** (1990) 576.
- [15] G. Dvali and M. Shifman, Nucl. Phys. **B504** (1997) 127; Phys. Lett. **B396** (1997) 64.
- [16] V. Kaplunovsky, Nucl. Phys. **B307** (1988) 145; (E) Nucl. Phys. **B382** (1992) 436.
- [17] C. Bachas and C. Fabre, Nucl. Phys. **B476** (1996) 418.
- [18] K.R. Dienes, Phys. Rept. **287** (1997) 447.
- [19] E. Cáceres, V.S. Kaplunovsky and I.M. Mandelberg, Nucl. Phys. **B493** (1997) 73.
- [20] S. Dimopoulos and H. Georgi, Nucl. Phys. **B150** (1981) 193;
W.J. Marciano and G. Senjanović, Phys. Rev. **D25** (1982) 3092;
C. Giunti, C.W. Kim and U.W. Lee, Mod. Phys. Lett. **A6** (1991) 1745;
J. Ellis, S. Kelley and D.V. Nanopoulos, Phys Lett. **B249** (1990) 441;
U. Amaldi, W. de Boer and H. Furstenuau, Phys. Lett. **B260** (1991) 447;
P. Langacker and M. Luo, Phys. Rev. **D44** (1991) 817.
- [21] I. Antoniadis, Phys. Lett. **B246** (1990) 377;
I. Antoniadis, C. Muñoz and M. Quirós, Nucl. Phys. **B397** (1993) 515;
I. Antoniadis, K. Benakli and M. Quirós, Phys. Lett. **B331** (1994) 313;
I. Antoniadis, S. Dimopoulos and G. Dvali, Nucl. Phys. **B516** (1998) 70.
- [22] T.R. Taylor and G. Veneziano, Phys. Lett. **B212** (1988) 147.
- [23] J. Polchinski, “TASI Lectures on D-Branes”, hep-th/9611050.
- [24] See, *e.g.*,
W. Lerche, Nucl. Phys. **B308** (1988) 102;
V. Kaplunovsky and J. Louis, Nucl. Phys. **B444** (1995) 191;
M. Bershadsky, S. Cecotti, H. Ooguri and C. Vafa, Commun. Math. Phys. **165** (1994) 311;
J.A. Harvey and G. Moore, Nucl. Phys. **B463** (1996) 215;
M.R. Douglas and M. Li, hep-th/9604041.
- [25] G. ’t Hooft, Nucl. Phys. **B72** (1974) 461.
- [26] M. Bershadsky, Z. Kakushadze and C. Vafa, Nucl. Phys. **B523** (1998) 59.

- [27] C.P. Bachas, JHEP **9811** (1998) 023.
- [28] Z. Kakushadze, Nucl. Phys. **B552** (1999) 3; Nucl. Phys. **B551** (1999) 549.
- [29] Z. Kakushadze, Phys. Lett. **B455** (1999) 120; hep-th/9904211; Phys. Lett. **B459** (1999) 497.
- [30] M.K. Gaillard and T.R. Taylor, Nucl. Phys. **B381** (1992) 577.
- [31] C.D. Carone, hep-ph/9902407.
- [32] A. Delgado and M. Quirós, hep-ph/9903400.